## Recommended COVID-19 vaccine doses<sup>1</sup>

### Primary course recommendations



6 months to 5 years 5 years to <18 18 years and older Not eligible Without risk factors **Recommended** a single primary dose. Eligible for 2 Eligible for a single primary doses and With risk factors<sup>2</sup> primary dose<sup>3</sup>. eligible for a 3rd primary dose<sup>3</sup>. With severe Immunocompromise **Recommended** 2 primary doses and **eligible** for a 3rd dose<sup>4</sup>.

#### Notes:


- 1. Monovalent Omicron XBB.1.5 vaccines are preferred; for ages in which a monovalent XBB.1.5-containing vaccine is not available, use other vaccines approved for that age group.
- 2. Includes those with a medical condition that increases the risk of severe COVID-19 illness (refer to <u>Australian Immunisation Handbook</u>) or those with disability with significant or complex health needs or multiple comorbidities which increase the risk of poor outcomes from COVID-19.
- 3. Consider dose based on an individual risk benefit assessment with an immunisation provider.
- 4. People with severe immunocompromise, who are over 6 months of age, are recommended 2 primary doses and are eligible for a 3<sup>rd</sup> primary dose based on an individual risk-benefit assessment. Please refer to the <u>Australian Immunisation Handbook</u> for further information.

Information current as of 1 March 2024.

# Recommended COVID-19 vaccine doses<sup>1</sup>



### COVID-19 Booster dose recommendation



#### Notes:

- 1. Monovalent Omicron XBB.1.5 vaccines are preferred; for ages in which a monovalent XBB.1.5-containing vaccine is not available, use other vaccines approved for that age group.
- 2. Consider dose based on an individual risk benefit assessment with an immunisation provider.
- 3. People with severe immunocompromise, who are aged between 18 to 64 years, are recommended a dose every 12 months and are eligible for a dose every 6 months based on an individual risk-benefit assessment. Please refer to the <u>Australian Immunisation Handbook</u> for further information.
- 4. People aged between 65 and 74 years are recommended a dose every 12 months and are eligible for a dose every 6 months based on an individual risk-benefit assessment. Please refer to the <u>Australian Immunisation Handbook</u> for further information.